

Integrated motions of molecular machines and motors: Small-angle scattering studies

Giacomo Mariani^a, Nicolas Jouault^a, Antoine Goujon^b, Jean-Rémy Colard-Itté^b, Emilie Moulin^b, Nicolas Giuseppone^b, and Eric Buhler^{a*}

- a. Matière et Systèmes Complexes Laboratory (MSC), UMR 7057, University Paris Diderot-Paris 7, Bâtiment Condorcet, 75205 Paris Cedex 13, France
- b. Institut Charles Sadron (ICS), University of Strasbourg, 23 rue du Loess, BP 84047, 67034 Strasbourg Cedex 2, France

* eric.buhler@univ-paris-diderot.fr

Switchable functional molecules capable of producing mechanical work constitute an active focus in nanotechnologies as they can be a source of components for molecular-based devices and materials. In particular, the dynamic nature of mechanically interlocked molecules allows their components to undergo relative internal movements, which can be exploited in translation and circumrotation. When it comes to using molecular machines to facilitate the creation of materials on the macro-scale, the primary concern is whether the nano-sized machines will be able to amplify their mechanical behavior to create a response in the bulk material. Hence, one of the most fundamental and challenging objectives associated to nano-machines rests on their coupling (in space and time) in order to transfer controlled motions from the molecular arena to the supramolecular and macroscopic scale.

In the present work, we have developed two kinds of responsive contractile polymeric materials: i) The first one concerns nano-machines linked into a supramolecular polymer in which we produced micrometric motions (contraction/extension) by the integration of thousands of single contractile nano-switches by altering the pH of the solution; just like myofibrils do when packed in bundles in muscles. ii) The second one is based on the connection of light-driven rotary motors with the possibility to make them functioning far from equilibrium and acting as reticulation units in an entangled polymer network.

Small-angle neutron scattering (coupled with light and X-ray scattering) has been used to investigate the structure of the supramolecular self-assemblies of nano-machines before and after the induced structural changes as well as the dynamics of the contraction process at different length and time scales. We discuss here the relation between the local and overall structure of the self-assemblies and the properties of the materials [1]. We show that these findings open up new possibilities of using molecular machines in smart responsive materials.

[1] G. Mariani, A. Goujon, E. Moulin, M. Rawiso, N. Giuseppone, E. Buhler *Nanoscale* **2017**, 9, 18456–18466, Integration of molecular machines into supramolecular materials: actuation between equilibrium polymers and crystal-like gels

